Important safety information for patients

Remember to have your liver function testing completed as instructed by your kidney doctor.

- **JINARC® can cause liver injury**. Liver function testing should be performed monthly for the first 18 months then every 3 months. Signs of liver injury include: nausea, vomiting, fever, tiredness, loss of appetite, pain in the abdomen, dark urine, yellowing of the skin or eyes, itching, joint and muscle pain.
- **JINARC®** should be stopped or discontinued if significant increase of liver enzymes and/or clinical symptoms of liver injury persist.
- **JINARC® can cause severe dehydration**. JINARC® blocks the action of vasopressin in the kidney and can cause large urinary volumes. It is recommended to drink plenty of water even if you do not feel thirsty to avoid excessive thirst or dehydration.
- **JINARC® must not be taken if you are pregnant or while breastfeeding.**

JINARC® is a registered trademark of Otsuka Pharmaceutical Co., Ltd. © Otsuka Australia Pharmaceutical Pty Ltd, ABN 20 601 768 754. Chatswood NSW 2067. Date of preparation: May 2026. JIN-AU-496.

Otsuka

My name:

Date JINARC® first prescribed:

My doctor's name:

Doctor's phone:

My pharmacy:

Pharmacy phone:

I am on **JINARC®** (tolvaptan),
a vasopressin antagonist.

For **JINARC®** Consumer Medicine Information
or Product Information call **1800 059 606**,
scan QR code or visit:

https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2018-CMI-01215-1